ACTEMRA® Infusion



Phone: 877.385.0535

Pleas	·	rferral form below and fax to OptiMed a cards, the patient's demographic face sl	•			
Prescriber Information	Prescriber:	NPI:				
		Fax: Office Contact:				
Patient Information		DOB: □M				
		Address:				
		2 nd Phone:				
	Primary Language:	Functional Limitations:				
	Diagnosis (include ICD-10 code):					
Clinical Information	Weight: □lb □kg Height: □in IV access: □PIV □PICC □Port □Other:					
	Patient's first dose of IV ACTEMRA®?					
	Allergies: Latex allergy? \(\sqrt{Y}\)es \(\sqrt{N}\)o					
	Prior treatments & reason for discontinuation:					
	Date of <u>negative</u> TB test: or □TB test pending, will fax results. Patient is HBV negative or has been treated: □Yes □No					
call	History of kidney disease: □Yes □No If yes, SCr: GFR/CrCl: History of heart failure: □Yes □No					
Clini	Required Labs: ANC	Platelets AST		ALT	SCr	
9	Result:	Result: (II	ı Ni·	Result: (ULN:		
	Date:	nesure	LIV,	Nesuit(OLIV	=/	
				1		
	Referring provider's preferred site of care*: OptiMed Infusion Center Home Infusion* OptiMed to determine site of care *Site of care preference is subject to payer limitations, clinical appropriateness, and the availability of servicing providers.					
	Based on the clinical judgement ACTEMRA® Dose	t of the pharmacist, doses may be roun Infusion Diluent/Volume	nded up or dow Rate	vn by no more than 10% unless c Frequency	hecked here: Number of Doses	
	Adult Rheumatoid Arthritis*	Intusion Diluent/Volume			Nullinet of Doses	
_	□4mg/kg	in 100ml NaCl 0 004	Infused over 60 minutes	every rour weeks		
ıtion	□8mg/kg <u>Polyarticular JIA</u>	in 100mL NaCl 0.9%				
rma	□10mg/kg (weight <30kg)	Weight <30kg: in 50mL NaCl 0.9%	Infused over	every rour weeks		
Info	□8mg/kg (weight ≥30kg) Systemic JIA	Weight ≥30kg: in 100mL NaCl 0.9%	60 minutes			
Prescription Informat	□12mg/kg (weight <30kg)	Weight <30kg: in 50mL NaCl 0.9%	Infused over	every two weeks		
	□8mg/kg (weight ≥30kg) *Doses exceeding 800mg per in	Weight ≥30kg: in 100mL NaCl 0.9% Ifusion are not recommended.	60 minutes			
	Premedication orders:					
	PRN medication orders:					
	Laboratory orders: □ANC/ Platelets/ AST/ ALT four to eight (4 to 8) weeks after the start of therapy and every three (3) months thereafter.					
	Other lab orders (subject to availability):					
		firms that the treatment(s) indicated on this referral				
Prescriber Signature		insurance prior authorization process and to provi			•	
, -	Signature: Date:					
esci						